

RAPID DUST PRODUCTION IN SUBMILLIMETER GALAXIES AT $Z > 4$?

MICHAŁ J. MICHAŁOWSKI^{1,2}, DARACH WATSON¹, AND JENS HJORTH¹

Astrophysical Journal 712 (2010) 942-950

ABSTRACT

The existence of submillimeter-selected galaxies (SMGs) at redshifts $z > 4$ has recently been confirmed. Using simultaneously all the available data from UV to radio we have modeled the spectral energy distributions of the six known spectroscopically confirmed SMGs at $z > 4$. We find that their star formation rates (average $\sim 2500 M_{\odot} \text{ yr}^{-1}$), stellar ($\sim 3.6 \times 10^{11} M_{\odot}$) and dust ($\sim 6.7 \times 10^8 M_{\odot}$) masses, extinction ($A_V \sim 2.2$ mag), and gas-to-dust ratios (~ 60) are within the ranges for $1.7 < z < 3.6$ SMGs. Our analysis suggests that infrared-to-radio luminosity ratios of SMGs do not change up to redshift ~ 5 and are lower by a factor of ~ 2.1 than the value corresponding to the local IR-radio correlation. However, we also find dissimilarities between $z > 4$ and lower-redshift SMGs. Those at $z > 4$ tend to be among the most star-forming, least massive and hottest (~ 60 K) SMGs and exhibit the highest fraction of stellar mass formed in the ongoing starburst ($\sim 45\%$). This indicates that at $z > 4$ we see earlier stages of evolution of submillimeter-bright galaxies. Using the derived properties for $z > 4$ SMGs we investigate the origin of dust at epochs less than 1.5 Gyr after the big bang. This is significant to our understanding of the evolution of the early universe. For three $z > 4$ SMGs, asymptotic giant branch stars could be the dominant dust producers. However, for the remaining three only supernovae (SNe) are efficient and fast enough to be responsible for dust production, though requiring a very high dust yield per SN ($0.15\text{--}0.65 M_{\odot}$). The required dust yields are lower if a top-heavy initial mass function or significant dust growth in the interstellar medium is assumed. We estimate lower limits of the contribution of SMGs to the cosmic star formation and stellar mass densities at $z \sim 4\text{--}5$ to be $\sim 4\%$ and $\sim 1\%$, respectively.

Subject headings: galaxies: active – galaxies: evolution – galaxies: high-redshift – galaxies: ISM – galaxies: starburst – submillimeter

1. INTRODUCTION

Submillimeter-selected galaxies (SMGs) are among the most powerful starburst galaxies in the Universe. Most of them have been found at redshifts 1.5–3 (Chapman et al. 2005). Their complex selection criteria (Blain et al. 2004), in particular the requirement of a radio detection to obtain a precise localization, make it difficult to discover the very high redshift tail of SMGs. This was addressed using deep, high resolution observations of SMGs (Iono et al. 2006; Tacconi et al. 2006, 2008; Wang et al. 2007, 2009; Younger et al. 2007, 2008a,b, 2009b,a; Dannerbauer et al. 2008; Cowie et al. 2009). The existence of SMGs at $z > 4$ has recently been spectroscopically confirmed by Coppin et al. (2009), Capak et al. (2008), Schinnerer et al. (2008), Daddi et al. (2009b,a), and Knudsen et al. (2008, 2010).

At these redshifts the age of the Universe is < 1.5 Gyr, which enforces the need for careful analysis of the timescales for formation of stars and dust. The important question is if supernovae (SNe), or asymptotic giant branch (AGB) stars, or some other sources are responsible for production of dust residing in these galaxies. Locally, dust is predominantly formed by evolved, post-main-sequence stars (Gehrz 1989), but the situation may be different at high redshifts. Dwek et al. (2007) claimed that only SNe can produce dust on timescales

less than 1 Gyr, but it has been shown by Valiante et al. (2009) that AGB stars begin to dominate dust production over SNe as early as 150–500 Myr after the onset of star formation (see also Sloan et al. 2009). SN-origin dust has been claimed to be present in $z \sim 6.2$ quasar (Maiolino et al. 2004) and in two gamma-ray burst host galaxies at $z \sim 6.3$ (Stratta et al. 2007, but see other interpretation of their data in Zafar et al. 2010) and at $z \sim 5$ (Perley et al. 2009).

In order to understand the formation of SMGs and their evolution through cosmic time, it is also important to compare high- and low-redshift SMG samples. This may help to constrain when their stars were formed.

SMGs at $z > 4$ are also suitable to study the infrared (IR) - radio correlation. This remarkably tight correlation, found locally (Helou et al. 1985; Condon 1992), was studied at redshifts $z \lesssim 3.5$ (Garrett 2002; Gruppioni et al. 2003; Appleton et al. 2004; Kovács et al. 2006; Boyle et al. 2007; Marleau et al. 2007; Vlahakis et al. 2007; Yang et al. 2007; Beswick et al. 2008; Hainline 2008; Ibar et al. 2008; Sajina et al. 2008; Garn et al. 2009; Michałowski et al. 2010; Murphy et al. 2009; Murphy 2009; Rieke et al. 2009; Seymour et al. 2009; Younger et al. 2009b; Sargent et al. 2010). No significant evolution of the correlation was found up to these redshifts, but SMGs seem to form a correlation by their own offset toward higher radio luminosities (Kovács et al. 2006; Michałowski et al. 2010; Murphy et al. 2009; Murphy 2009). The only sign of evolution was reported by Ivison et al. (2010) based on stacking analysis of 24 μm -selected galaxies, though possibly interpreted as a

¹ Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen Ø, Denmark; michal@dark-cosmology.dk

² Scottish Universities Physics Alliance, Institute for Astronomy, University of Edinburgh, Royal Observatory, Edinburgh, EH9 3HJ, UK

TABLE 1
PHOTOMETRY DETECTIONS OF $z > 4$ SMGs

SMG	z	λ_{obs} (μm)	Flux (μJy)	Error (μJy)	Ref.
LESSJ033229.4-275619	4.760	0.597	0.072	0.006	Coppin09
LESSJ033229.4-275619	4.760	0.652	0.215	0.020	Coppin09
LESSJ033229.4-275619	4.760	0.771	0.350	0.013	Coppin09
LESSJ033229.4-275619	4.760	0.905	0.405	0.015	Coppin09
LESSJ033229.4-275619	4.760	3.600	2.860	0.060	Coppin09

NOTE. — The measured fluxes (Column 4) are given with the corresponding 1σ errors (Column 5). The data for SMMJ163555.5+661300 have been corrected for lensing magnification of a factor of 5.5. (This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.)

REFERENCES. — Capak et al. (2004, 2008); Giavalisco et al. (2004); Wang et al. (2004, 2007, 2009); Frayer et al. (2006); Iono et al. (2006); Pope et al. (2006); Dannerbauer et al. (2008); Greve et al. (2008); Perera et al. (2008); Schinnerer et al. (2008); Casey et al. (2009); Chapin et al. (2009); Coppin et al. (2009); Daddi et al. (2009b,a); Devlin et al. (2009); Knudsen et al. (2010)

TABLE 2
PHOTOMETRY UPPER LIMITS OF $z > 4$ SMGs

SMG	z	λ_{obs} (μm)	Flux (μJy)	Error (μJy)	Ref.
LESSJ033229.4-275619	4.760	0.352	0.087	0.000	Coppin09
LESSJ033229.4-275619	4.760	0.433	0.013	0.000	Coppin09
LESSJ033229.4-275619	4.760	0.461	0.039	0.000	Coppin09
LESSJ033229.4-275619	4.760	0.538	0.059	0.000	Coppin09
SMMJ123711.7+622212	4.055	0.365	0.063	0.026	Capak04

NOTE. — When the error is equal to zero, the flux column denotes the 3σ upper limit. Otherwise it denotes formal flux at the position of an SMG. The data for SMMJ163555.5+661300 have been corrected for lensing magnification of a factor of 5.5. (This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.)

REFERENCES. — Capak et al. (2004, 2008); Giavalisco et al. (2004); Wang et al. (2004, 2007, 2009); Frayer et al. (2006); Iono et al. (2006); Pope et al. (2006); Dannerbauer et al. (2008); Greve et al. (2008); Perera et al. (2008); Schinnerer et al. (2008); Casey et al. (2009); Chapin et al. (2009); Coppin et al. (2009); Daddi et al. (2009b,a); Devlin et al. (2009); Knudsen et al. (2010)

selection effect. It is however possible that the correlation breaks down at even earlier epochs due to changes in star formation processes, e.g., suppression of radio emission in inverse Compton losses off the CMB photons as suggested by Lacki et al. (2009), Lacki & Thompson (2009), and Murphy (2009).

In Michałowski et al. (2010) we analyzed the full UV-to-radio spectral energy distributions (SEDs) of 76 SMGs from the Chapman et al. (2005) sample with spectroscopic redshifts up to $z < 3.6$. Here we extend that study by analyzing the sample of all spectroscopically confirmed SMGs at $z > 4$. The main objective of this paper is to characterize the required efficiency of dust producers (SNe and AGB stars) at these early epochs of the evolution of the Universe. In Section 2 the SMG sample is presented. We outline our methodology and derive the properties of SMGs in Section 3 and discuss the implications in Section 4. Section 5 closes with our conclusions. We use a cosmological model with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.7$ and $\Omega_m = 0.3$.

2. SAMPLE

We selected all six SMGs with robust (optical or CO) redshifts at $z > 4$ identified by Coppin et al. (2009), Capak et al. (2008), Schinnerer et al. (2008), Daddi et al. (2009b,a), and Knudsen et al. (2008, 2010) in the ECDF-S (900 arcmin²; Coppin et al. 2009; Greve et al. 2009; Weiß et al. 2009), COSMOS (1080 arcmin²; Scott et al. 2008), GOODS-N (100 arcmin²; Hughes et al. 1998; Barger et al. 2000; Chapman et al. 2001; Borys et al. 2003; Serjeant et al. 2003; Wang et al. 2004), and Abell 2218 (11.8 arcmin²; Knudsen et al. 2006) fields. The photometric data are presented in Tables 1 and 2. The data for SMMJ163555.5+661300 have been corrected for lensing magnification of a factor of 5.5 obtained by Knudsen et al. (2010) using the model of Elíasdóttir et al. (2007).

Our sample is not homogeneously selected. Namely, some of the sources are bright enough in the optical to allow spectroscopy, whereas redshifts of some of them were measured based on CO emission (one even not detected at optical wavelengths). In particular, LESSJ033229.4-275619 and SMMJ100054.48+023435.9 were targeted spectroscopically, because they are V-dropouts suggesting $z \approx 5$ (Coppin et al. 2009; Capak et al. 2008) and their optical counterparts (and hence, redshifts) are based on detections at radio wavelengths, which likely biases them toward high star formation rates (SFRs). The CO lines from SMMJ123711.7+622212 (called GN20 in Pope et al. 2006) and SMMJ123709.5+622206 (GN20.2a) were detected serendipitously while observing an angularly close galaxy at $z = 1.522$ (Daddi et al. 2009b), and the CO emission of SMMJ123633.8+621408 (GN10) was searched for under the assumption that it is a member of the protocluster structure containing GN20 and GN20.2a (Daddi et al. 2009a). Finally, SMMJ163555.5+661300 was detected because of its lensing magnification and is therefore intrinsically the faintest member of our sample.

It is therefore not easy to quantify how selection effects influence our results. Very likely our sample is biased toward high luminosity objects, i.e., with high SFRs. This is supported by the fact that SMMJ163555.5+661300, magnified by lensing and the only member of our sample not detected in the radio, has a much lower SFR than blank-field members of our sample (Table 3). However, given the significance of the *Spitzer* IRAC detections (rest-frame $\sim 1\text{--}2 \mu\text{m}$) of $\gtrsim 10\sigma$, our sample is *not* biased against low stellar masses.

3. SED FITTING AND RESULTS

We applied the SED fitting method detailed in Michałowski et al. (2008, 2009, 2010, see therein a discussion of the derivation of galaxy properties and typical uncertainties) based on 35 000 templates in the library of Iglesias-Páramo et al. (2007), plus some templates of Silva et al. (1998) and Michałowski et al. (2008), all developed in GRASIL (Silva et al. 1998). The templates cover a broad range of galaxy properties and were tested to reproduce the SEDs of high-redshift galaxies (Silva et al. 1998; Iglesias-Páramo et al. 2007; Michałowski et al. 2008, 2010). Their star formation histories are assumed to be a smooth Schmidt-type law (SFR proportional to the gas mass to some power, see Silva et al. 1998, for details) with a starburst (if any) on top of that starting 50 Myr before the time at which




FIG. 1.— Spectral energy distributions (SEDs) of $z > 4$ SMGs. *Solid lines*: the best GRASIL fits. *Squares*: detections with errors, in most cases, smaller than the size of the symbols. *Arrows*: 3σ upper limits (values marked at the base). The data for SMMJ163555.5+661300 have been corrected for lensing magnification of a factor of 5.5.

TABLE 3
PROPERTIES OF THE $z > 4$ SMGS DERIVED FROM THE SED MODELING

SMG (1)	z (2)	SFR ($M_{\odot} \text{ yr}^{-1}$)			SSFR (Gyr^{-1}) (7)	M_* ($10^{11} M_{\odot}$) (8)	M_{burst}/M_* (%) (9)	M_*/L_K ($M_{\odot}/L_{\odot}$) (10)	M_d ($10^9 M_{\odot}$) (11)	L_{IR} ($10^{13} L_{\odot}$) (12)	T_d (K) (13)	A_V (mag) (14)	q (15)	AGN? (16)
		SED (3)	UV (4)	IR (5)										
LESSJ033229.4-275619	4.760	1718	26	1150	1636	1.82	6.3	12.5	0.7	0.7	38.7	2.01	2.36	SB,spec
MMJ100054.48+023435.9	4.547	3316	60	2725	3548	1.438	1.9	82.3	0.4	1.6	71.5	1.77	2.40	SB
SMMJ123711.7+622212(GN20,AzGN01)	4.055	6120	33	4577	4498	5.61	8.2	33.5	1.1	2.7	61.3	2.55	2.52	...
SMMJ123709.5+622206(GN20.2a,AzGN1.2)	4.051	891	27	841	10700	3.44	2.4	14.3	1.4	0.5	38.7	1.76	1.41	rad
SMMJ123633.8+621408(GN10,GN850-5,AzGN03)	4.042	2877	0	5031	1951	28.57	1.8	100.0	0.3	2.9	97.2	7.84	2.93	SB
SMMJ163555.5+661300	4.044	339	1	289	<483	3.40	0.9	18.1	0.2	0.2	45.1	2.91	>2.29	SB

NOTE. — Column (1): SMG name (alternative names from Wang et al. 2004, 2009; Pope et al. 2006; Daddi et al. 2009b,a; Perera et al. 2008; Chapin et al. 2009). Column (2): redshift (Coppin et al. 2009; Capak et al. 2008; Daddi et al. 2009b,a; Knudsen et al. 2010). Column (3): total star formation rate (SFR) for 0.15 – 120 $M_{\odot}$ stars averaged over the last 50 Myr derived from the SED model. Column (4): SFR from UV emission interpolated from the SED template (using Kennicutt 1998). Column (5): SFR from IR emission (Column 12) used in all analysis throughout the paper (using Kennicutt 1998). Column (6): SFR from radio emission derived directly from the radio data (using Bell 2003). Column (7): specific SFR $\equiv \text{SFR}_{\text{IR}}/M_*$. Column (8): stellar mass. Column (9): Ratio of the mass of gas converted to star during the recent starburst episode to the total stellar mass. Column (10): stellar mass to light ratio (luminosity at rest-frame K was interpolated using the best SED model). Column (11): total 8 – 1000 μm infrared luminosity. Column (13): dust temperature (we assumed emissivity index $\beta = 1.3$). Column (14): Average extinction $A_V = 2.5 \log(V\text{-band starlight unextinguished} / V\text{-band starlight observed})$. Column (15): IR-radio correlation parameter (Section 4.3). Column (16): AGN flag. — SB: X-ray identified starburst (no X-ray detection; Coppin et al. 2009; Capak et al. 2008; Dannerbauer et al. 2008; Wang et al. 2009; Knudsen et al. 2010), as opposed to X-ray identified AGN (Alexander et al. 2005); spec: spectroscopically identified AGN (Coppin et al. 2009); rad: radio datapoint is more than 3σ above the starburst model.

the SED is computed. There are seven free parameters in the library of Iglesias-Páramo et al. (2007): the normalization of the Schmidt-type law, the timescale of the mass infall, the intensity of the starburst, the timescale of the molecular cloud destruction, the optical depth of molecular clouds, the age of a galaxy and the inclination of a disk with respect to the observer.

We scaled all the SEDs to match the data (detections) and chose the one with the lowest χ^2 . In all but one case (see below) the upper limits did not provide a constraint on the best-fitting model.

In all but one case we obtained the best fits using the library of Iglesias-Páramo et al. (2007). For GN10 we fitted a model corresponding to a 0.1 Gyr old progenitor of an elliptical galaxy (Silva et al. 1998) with modified maximum grain temperatures from 400 K to 100 K in order to suppress strong mid-IR emission in the original model, otherwise inconsistent with the data. All other models failed to reproduce its extremely red observed-frame 2.1–3.6 μm color (Wang et al. 2009).

The best fits³ are shown in Figure 1. It is apparent that GRASIL models can reproduce the SEDs of even such distant galaxies. The resulting properties of the galaxies are listed in Table 3.

In order to assess the uncertainties of the derived parameters, we calculated the confidence intervals using SED models with $\Delta\chi^2 < 1$. The resulting uncertainties are a factor of ~ 2 –4 for SFRs and L_{IR} ; a factor of ~ 1.5 –2.5 for M_* ; a factor of ~ 2 –5 for M_{burst} ; ~ 10 K for T_d ; and 0.3–1.0 mag for A_V . The uncertainty on T_d introduces a factor of ~ 1.5 –2.5 error on M_d and allowing the emissivity β to vary in the range 1–2 results in an uncertainty of a factor of < 3 on M_d . The choice of a Salpeter (1955) initial mass function (IMF) with cutoffs of 0.15 and 120 $M_{\odot}$ introduces a maximum systematic error of a factor of ~ 2 in the determination of the stellar masses and SFRs (Erb et al. 2006).

Our estimates of SFRs, L_{IR} and M_d are consistent within a factor of < 3 with those obtained by Coppin et al. (2009), Capak et al. (2008), Schinnerer et al. (2008), Younger et al. (2008b), Casey et al. (2009), Daddi et al. (2009b,a), Knudsen et al. (2010), and Wang et al. (2009) (compare Tables 3 and 4). Our M_* estimates are consistent with those of Daddi et al. (2009b,a), and Wang et al. (2009) after taking into account that Daddi et al. (2009b,a) used a Chabrier (2003) IMF resulting in stellar masses 1.8 times lower than for the Salpeter (1955) IMF (Erb et al. 2006).

However we obtained stellar masses for LESSJ033229.4-275619 and MMJ100054.48+023435.9 ~ 10 times larger than Coppin et al. (2009)⁴ and Capak et al. (2008), respectively, and for SMMJ163555.5+661300 ~ 5 times larger than Knudsen et al. (2010). The difference can be explained by the fact that Coppin et al. (2009) assumed a mass-to-light ratio of $M_*/L_K = 0.1$ and the determinations of Capak et al. (2008) and Knudsen et al. (2010) correspond to $M_*/L_K \sim 0.03$ and ~ 0.17 , respectively. These are very low values compared to lower-

³ The SED fits can be downloaded from <http://archive.dark-cosmology.dk>

⁴ Note that our estimate agrees with that of Stark et al. (2007).

TABLE 4
PROPERTIES OF $z > 4$ SMGS DERIVED BY OTHER AUTHORS

SMG	SFR ($M_{\odot} \text{ yr}^{-1}$)	L_{IR} ($10^{13} L_{\odot}$)	M_{*} ($10^{11} M_{\odot}$)	M_d ($10^9 M_{\odot}$)	Ref.
LESS	1000	0.61	0.5	0.5	Coppin et al. (2009)
MM	1000-4000	0.5–2.0	0.1		Capak et al. (2008), Schinnerer et al. (2008)
GN20		1.0–5.0	2.3		Daddi et al. (2009b), Younger et al. (2008b), Casey et al. (2009)
GN20.2a		0.5–1.6	0.5		Daddi et al. (2009b), Casey et al. (2009)
GN10	2400	1.2–2.5	1.0–3.0		Daddi et al. (2009a), Wang et al. (2009)
SMM	230	0.13	0.16		Knudsen et al. (2010)

NOTE. — The first column lists all SMGs in the order given in Table 3. Only the first parts of their names are given for brevity.

redshift galaxies (e.g., Drory et al. 2004; Portinari et al. 2004; Labbé et al. 2005; Castro Cerón et al. 2006, 2009; van der Wel et al. 2006; Courty et al. 2007; Michałowski et al. 2010; Savaglio et al. 2009) giving lower limits on stellar masses. On the other hand, we do not assume mass-to-light ratios, but derive them from the stellar population models incorporated in GRASIL.

4. DISCUSSION

4.1. Formation of Stars in $z > 4$ SMGs

All $z > 4$ SMGs in our sample are characterized by an extremely strong starburst episode (average SFR $\sim 2500 M_{\odot} \text{ yr}^{-1}$, Column 5 of Table 3) during which a substantial fraction (average $\sim 45\%$, Column 9) of their stellar population was formed. They are therefore manifestations of the strongest known star-forming events in the universe. Their high stellar masses (average $\sim 3.6 \times 10^{11} M_{\odot}$) agree with a suggestion of Davé et al. (2009) based on numerical simulations that the most rapidly star forming galaxies coincide with the most massive galaxies.

The properties of the galaxies are within the ranges found by Michałowski et al. (2010) for the $1.7 < z < 3.6$ SMG sample (Chapman et al. 2005). However, $z > 4$ SMGs tend to be among the most star-forming, least massive, and hottest SMGs and exhibit the highest fraction of stellar mass formed in the ongoing starburst. Namely, 43% of $1.7 < z < 3.6$ SMGs have lower SFRs than any of the $z > 4$ SMGs⁵, whereas only 2% have higher SFRs; 30% of $1.7 < z < 3.6$ SMGs have higher stellar masses, whereas only 8% have lower stellar masses; 55% have a lower fraction of stellar mass formed in the ongoing starburst episode, whereas only 2% have higher fraction; and 28% have lower dust temperatures, whereas only 4% have higher temperature. This can be interpreted as SMGs at $z > 4$ representing earlier stages of the formation of submillimeter-bright galaxies at which the pre-existing stellar population is less pronounced and therefore the ongoing starburst episode contributes more to the final stellar mass.

However, we note that we cannot exclude the possibility that these galaxies are AGN dominated, suggested by their high IR luminosities and dust temperatures. In that case the SFRs and M_{*} we derive would be upper limits.

If GN10 is not AGN dominated, then the fact that we could only fit its SED using a template corresponding to a young (0.1 Gyr old) progenitor of an elliptical

galaxy supports the hypothesis that SMGs evolve into ellipticals. According to our model, after 1.5 Gyr GN10 will evolve into a massive elliptical containing $\sim 10^{12} M_{\odot}$ stars. In order to explain its extremely red observed-frame 2.1–3.6 μm color, we did not need to invoke an old stellar population as suggested by Wang et al. (2009). Its spectrum is reproduced by a young stellar population residing in molecular clouds heavily obscured by dust with $A_V \sim 7.8$ mag (Table 3, Column 14). This estimate is similar to the values obtained using only near-IR and optical data by Daddi et al. (2009a) and Wang et al. (2009).

On the other hand the remaining $z > 4$ SMGs are only moderately obscured ($A_V \sim 2$ mag; Column 14). We checked that this value of extinction is roughly consistent with the expectation based on the estimation of the amount of energy reprocessed by dust. A crude estimate of the UV (0.2 μm) extinction can be calculated using $A_{0.2 \mu\text{m}} = 2.5 \log(\text{SFR}_{\text{IR}}/\text{SFR}_{\text{UV}})$. This gives $A_{0.2 \mu\text{m}} \sim 20$ mag for GN10 and ~ 3.8 –6.3 mag for the other objects. Assuming that $A_{0.2 \mu\text{m}} \sim 3 \times A_V$ (e.g., Gordon et al. 2003), this corresponds to $A_V \sim 6.6$ mag for GN10 and $A_V \sim 1.2$ –2.1 mag for the other objects. These values are consistent with those derived from the SED modeling (Table 3).

We estimated the SMG comoving volume densities as a sum of SFRs (or L_{IR} , or stellar or dust masses) of $z > 4$ SMGs⁶ divided by the comoving volume in the redshift interval 4–5 ($5.9 \times 10^6 \text{ Mpc}^3$). We obtained $\rho_{\text{SFR}} = 2.6 \times 10^{-3} M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$ (a similar value was found by Younger et al. 2009a, based on candidates for high-redshift SMGs), $\log \rho_{L_{\text{IR}}} = 7.17 L_{\odot} \text{ Mpc}^{-3}$, $\log \rho_{*} = 5.54 M_{\odot} \text{ Mpc}^{-3}$, and $\log \rho_{\text{dust}} = 2.81 M_{\odot} \text{ Mpc}^{-3}$. However one must keep in mind that these numbers could be affected by cosmic variance, because three out of five SMGs considered here are members of a protocluster structure (Daddi et al. 2009b,a). The contributions to the cosmic SFR and stellar mass densities of SMGs at these redshifts are 4.4% and 1.0%, respectively, using the compilation of the total values in Michałowski et al. (2010, Tables A.4 and A.5). These numbers indicate that currently detected SMGs did not contribute significantly to the cosmic star formation history at $z > 4$, but our estimates should be regarded as lower limits since more of such distant SMGs could still be undetected in the fields discussed here.

4.2. Producers of Dust in $z > 4$ SMGs

The dust masses we find for $z > 4$ SMGs (Table 3, Column 11) are similar to those derived for

⁵ We exclude SMMJ163555.5+661300 from this analysis, because its unlensed submillimeter flux of $\sim 2 \text{ mJy}$ makes it impossible to be detected by SCUBA in the blank-field survey similar to those used by Chapman et al. (2005).

⁶ We again exclude SMMJ163555.5+661300 because it is lensed.

TABLE 5
DUST YIELDS PER STAR REQUIRED TO EXPLAIN DUST IN $z > 4$ SMGS

Dust Producer	IMF	Total Mass	Dust Yields ($M_{\odot}$ Per Star)					
			LESS	MM	GN20	GN20.2a	GN10	SMM
AGB ($2.5 < M < 8M_{\odot}$)	Salpeter	M_*	0.03	0.06	0.04	0.17	0.05	0.07
AGB ($2.5 < M < 8M_{\odot}$)	Salpeter	$M_* - M_{\text{burst}}$	0.04	0.31	0.06	0.20	∞	0.09
SN ($M > 8M_{\odot}$)	Salpeter	M_*	0.14	0.22	0.15	0.65	0.18	0.27
SN ($M > 8M_{\odot}$)	Top-heavy	M_*	0.05	0.08	0.05	0.23	0.06	0.10

NOTE. — The IMF is either Salpeter (1955) with $\alpha = 2.35$ or top-heavy with $\alpha = 1.5$. The total mass indicates if the entire stellar mass (M_*) was used to calculate the number of stars (Equation A1), or if stars created during the ongoing starburst were excluded ($M_* - M_{\text{burst}}$). The last six columns contain dust yields for all SMGs in the order given in Table 3. Only the first parts of their names are given for brevity.

$z \sim 4$ –6 quasars (a few $\times 10^8 M_{\odot}$; Dunlop et al. 1994; Benford et al. 1999; Archibald et al. 2001; Omont et al. 2001; Priddey & McMahon 2001; Priddey et al. 2003, 2008; Isaak et al. 2002; Bertoldi et al. 2003a; Robson et al. 2004; Beelen et al. 2006; Wang et al. 2008; Martínez-Sansigre et al. 2009). These huge dust masses indicate that dust in SMGs was efficiently formed and able to survive even when the age of the universe was only 1.2–1.5 Gyr.

As detailed below we find that *i*) AGB stars are efficient and fast enough to form dust in the LESSJ033229.4-275619, GN20 and SMMJ163555.5+661300; *ii*) only SNe are efficient and fast enough to form dust in MMJ100054.48+023435.9, GN20.2a, and GN10, as long as the unusually high dust yields derived for Cassiopeia A and Kepler SN remnants are correct and typical, or if a top-heavy IMF and/or significant dust growth in the interstellar medium (ISM) is assumed. If these assumptions are correct, then SNe could also produce dust in SMGs mentioned in *i*.

These conclusions are not significantly affected by the uncertainties reported in Section 3. Even if dust masses were higher or stellar masses lower by a factor of a few, the required dust yields per SN would not exceed that of Cassiopeia A (Dunne et al. 2009) and Kepler (Gomez et al. 2009). It would however be more critical for AGB stars if dust masses were higher than we derived, since the required dust yields for LESSJ033229.4-275619, GN20, and SMMJ163555.5+661300 are already on the high end of the theoretically allowed yields. However, a decrease of dust masses by a factor of a few would not change our conclusion that AGB stars are not effective enough to form dust in MMJ100054.48+023435.9, GN20.2a, and GN10.

4.2.1. Asymptotic giant branch stars

In order to investigate whether AGB stars can be responsible for dust production in $z > 4$ SMGs, we estimated (see the Appendix) the average dust yields required per star with mass $2.5 < M < 8 M_{\odot}$ and main-sequence lifetime in a range 1 Gyr–55 Myr (calculated as $10^{10} \text{ yr} \times [M/M_{\odot}]^{-2.5}$; e.g., Kippenhahn & Weigert 1990). The lower mass limit was chosen to ensure that the stars considered can start producing dust within the age of the universe at the redshifts of our sources (1.2–1.5 Gyr).

The results are listed in the first row of Table 5. They are independent of the assumed star formation history of galaxies, but depend only on derived dust and stellar masses, assumed IMF and measured redshifts. We find that each AGB star would need to produce ~ 0.03 – $0.07 M_{\odot}$ of dust in order to explain the dust in $z > 4$

SMGs, excluding GN20.2a (see below). These numbers are close to the highest theoretical dust yields of AGB stars (Morgan & Edmunds 2003; Ferrarotti & Gail 2006, the total dust yields per AGB star are approximately metallicity independent) making them plausible dust producers. Assuming a top-heavy IMF does not change this result significantly (the required dust yields increase by $\sim 30\%$), because the choice of the slope of the IMF affects the number of high- and low-mass stars, leaving the number of intermediate-mass stars approximately constant. For GN20.2a the required dust yield ($0.17 M_{\odot}$) is too high to claim that AGB stars formed its dust.

However, MMJ100054.48+023435.9 and GN10 formed the majority of their stars in the ongoing starburst episode (Column 9, Table 3), which is too short for the $2.5 < M < 8 M_{\odot}$ stars considered above to finish their main-sequence phase. Therefore AGB stars could not contribute to the dust production in these two galaxies. To quantify this, we calculated the required dust yields for AGB stars taking into account only stars that were born before the ongoing starburst (replacing M_* by $M_* - M_{\text{burst}}$ in Equation A1). The resulting yields (the second row of Table 5) for LESSJ033229.4-275619, GN20, GN20.2a, and SMMJ163555.5+661300 do not differ significantly from our previous estimate (the first row of Table 5), because in this way we removed $\lesssim 30\%$ of stars (those formed during the ongoing starburst). However the yields for MMJ100054.48+023435.9 and GN10 become too high to claim that AGB stars formed dust in these galaxies.

A potential limitation of this claim is the uncertainty in determining the fraction of stellar mass formed during the ongoing starburst episode. If more stars were formed in the past, then AGB stars could be responsible for dust production in these two galaxies. However, even if it was the case, then the current SFRs would be the same as we derive, because they are fixed by strong submillimeter emission. Then the ongoing starburst becomes unfeasibly short (< 25 Myr) in order not to produce more stars than is inferred from the optical to near-infrared part of the spectra. Therefore, more stars in these two SMGs could have been formed before the ongoing starburst only if the current SFRs are overestimated due to a significant AGN contribution.

4.2.2. Supernovae

We repeated the analysis for SNe, i.e., calculated the required dust yields per one massive star ($> 8 M_{\odot}$). The yields are ~ 0.15 – $0.65 M_{\odot}$ of dust per SN (the third row of Table 5), consistent with the theoretical works (though without dust grain destruction imple-




FIG. 2.— Ratio of the infrared (8-1000 μm) and radio luminosities $q = \log(L_{\text{IR}}/3.75 \times 10^{12}/I_{1.4})$ as a function of redshift of SMGs. Both SMGs at $z > 4$ (red circles) and at lower redshifts (yellow, green, blue, violet circles, divided into four redshift bins) from Michałowski et al. (2010, M09) are shown. In the redshift range $1.4 < z < 5.0$ no significant evolution of the IR-radio correlation is found for SMGs. The average q for SMGs (squares) is however offset toward higher radio luminosities (factor of ~ 2.1 – 2.3) from the local relation ($q = 2.64$: solid line with scatter, 0.26: dotted lines; Bell 2003). A linear fit to the data (dashed line) resulted in q decreasing with redshift, but only at $\sim 2\sigma$ significance.

mented) of Todini & Ferrara (2001) and Nozawa et al. (2003); with a value predicted by Dwek et al. (2007) to account for dust in a $z \sim 6.4$ quasar; and with submillimeter observation of SN remnants Cassiopeia A (Dunne et al. 2003, 2009) and Kepler (Morgan et al. 2003; Gomez et al. 2009). There is a debate about the latter results on Cassiopeia A and Kepler (e.g., Dwek 2004; Krause et al. 2004; Gomez et al. 2005; Wilson & Batrla 2005; Blair et al. 2007; Sibthorpe et al. 2009), but if the dust yields of Dunne et al. (2009) and Gomez et al. (2009) were correct and typical, then SNe would be efficient enough to account for the dust in all $z > 4$ SMGs.

However our, required estimates are at least an order of magnitude higher than any other observed SN dust yields, which are typically in the range $\sim 10^{-3}$ – $10^{-2} M_{\odot}$ (Green et al. 2004; Borkowski et al. 2006; Sugerman et al. 2006; Ercolano et al. 2007; Meikle et al. 2007; Rho et al. 2008, 2009; Kotak et al. 2009; Sakon et al. 2009; Sandstrom et al. 2009; Wesson et al. 2009) and theoretically predicted dust masses able to survive in SN remnants (Bianchi & Schneider 2007). This apparent difficulty in explaining dust production in $z > 4$ SMGs can be resolved with a combination of two plausible effects.

Approximately half of the discrepancy can be accounted for with a top-heavy IMF giving more SNe per unit stellar mass (both top-heavy and Salpeter IMFs

have been claimed to reproduce the number counts of SMGs; Baugh et al. 2005; Fontanot et al. 2007). Changing the IMF slope from $\alpha = 2.35$ to $\alpha = 1.5$, consistent with values for low-mass star clusters (Scalo 1998) and a limit derived for a proto-star cluster (Sternberg 1998), resulted in the required dust yield decreasing to 0.05–0.23 $M_{\odot}$ (the fourth row of Table 5).

The second possibility is that SNe provided only the dust seeds and that the bulk of the dust mass was accumulated during grain growth in the ISM (e.g., Draine 2003). The timescale of this process is typically less than a few $\times 10$ Myr (Hirashita 2000; Zhukovska et al. 2008; Draine 2009), i.e., short enough to contribute significantly to the growth of dust mass in $z > 4$ SMGs. However, the extreme environments of $z > 4$ SMGs may make it difficult for grains to grow before they are destroyed. The hypothesis of dust growth can be tested by investigation whether the grains formed by SN remnants are smaller than those present in $z > 4$ SMGs.

4.3. The IR-Radio Correlation at $z > 4$

Despite the differences compared to the $1.7 < z < 3.6$ population (Section 4.1), the mean IR-to-radio luminosity ratio ($q \equiv \log[L_{\text{IR}}/3.75 \times 10^{12}/I_{1.4}]$ with radio K -corrections assuming a slope of -0.75) for $z > 4$ SMGs of 2.32 ± 0.20 is consistent with the values derived for lower-redshift SMGs (Michałowski et al. 2010). At $z > 4$ we find an offset of $\Delta q \sim -0.32$ (a factor of ~ 2.1) from

the local value of $q = 2.64$ (Bell 2003), though due to the small sample, this offset is significant only at $\sim 1.6\sigma$ level.

Hence, with our multi-wavelength approach we confirm the results of Murphy (2009), who derived the mean $q = 2.16 \pm 0.28$ for $z > 4$ SMGs. Moreover, our estimate is consistent with a value of 2.41 ± 0.20 derived for 250 μm -selected galaxies at $z < 3$ by Ivison et al. (2010).

The offset of $\Delta q \sim -0.32$ is consistent with a hypothesis of Lacki et al. (2009) and Lacki & Thompson (2009) that SMGs are “puffy starbursts” (vertically and radially extended galaxies with vertical scale heights ~ 1 kpc) experiencing weaker bremsstrahlung and ionization losses resulting in stronger radio emission. We cannot however exclude an AGN contribution boosting their radio fluxes. Since the redshifts of four out of six $z > 4$ SMGs have been measured independently of radio detections, the radio excess cannot be a result of a bias against radio-faint sources.

In Figure 2, we show the q -values as a function of redshift for both $z > 4$ SMGs discussed in this paper (red circles) and lower-redshift SMGs from Michałowski et al. (2010). The IR-radio correlation of SMGs does not show any significant evolution in the redshift range $1.4 < z < 5.0$. A linear fit to all the data results in a low-significance ($\sim 2\sigma$) indication that q decreases with redshift (as also found by Ivison et al. 2010, from a stacking analysis of 24 μm -selected galaxies) in the form $q = (-0.078 \pm 0.038)z + (2.50 \pm 0.09)$. To date, there are only two q determinations at higher redshifts than presented here, namely $z = 6.2$ and 6.42 quasars (Carilli et al. 2004; Beelen et al. 2006, $q = 1.8$ – 2.2).

4.4. Gas-to-Dust Ratio at $z > 4$

Using the molecular gas mass estimates (based on CO[4–3] line observations) from Schinnerer et al. (2008) and Daddi et al. (2009b,a) we derive gas-to-dust ratios of $M_{H_2}/M_d = 73, 47, 22$, and 98 for MMJ100054.48+023435.9, GN20, GN20.2a, and GN10, respectively. We note that these values are affected by significant uncertainties of the conversion from CO line strength to H_2 mass. The mean value of 60 is consistent with 54^{+14}_{-11} estimated for $z \sim 1$ – 3.5 SMGs by Kovács et al. (2006) using the CO survey of Greve et al. (2005). SMGs at $z > 4$ have one of the lowest gas-to-dust ratios compared to galaxies selected at other wavelengths, e.g., the Milky Way (~ 90 – 400 ; Sodroski et al. 1997), other spirals ($\sim 1000 \pm 500$; Devereux & Young 1990; Stevens et al. 2005), the nuclear regions of local luminous IR galaxies, ultraluminous IR galaxies (120 ± 28 ; Wilson et al. 2008), and of local, far-IR-selected galaxies (~ 50 ; Seaquist et al. 2004) (all these results are based on IR and CO data, and are therefore directly comparable to our estimates). This is not surprising, since SMGs are

selected by submillimeter emission. Similarly, a low value of $M_{H_2}/M_d = 30$ was found by Bertoldi et al. (2003b) for $z = 6.42$ quasar, but Cox et al. (2002) reported a higher value of $M_{H_2}/M_d = 150$ for $z \sim 4$ quasar. The small number of SMGs and quasars with derived gas-to-dust ratios hampers a comparison of these two samples.

5. CONCLUSIONS

We have analyzed the SEDs of six spectroscopically confirmed $z > 4$ SMGs. Our results provide constraints on dust production at these early epochs of the evolution of the universe. We find that AGB stars are efficient and fast enough to form the dust residing in three of these galaxies. However, for the three remaining SMGs only SNe are efficient and fast enough. The high required SN dust yields hint at a possibility that their stars may be distributed according to a top-heavy IMF and/or that dust grains are substantially grown in the ISM. Since the majority of the stars in these galaxies were formed on very short timescales, these properties are very likely to be similar to those of the first galaxies beyond redshift 6, which had been building up their stellar populations only for several hundred Myr, which elapsed since the big bang.

We present evidence that the IR-to-radio luminosity ratios of $z > 4$ SMGs are consistent with that of lower-redshift SMGs and are offset from the local relation by a factor of ~ 2.1 .

A comparison of the $z > 4$ SMGs with the lower-redshift sample, in particular their high SFRs, dust temperatures and fraction of stars formed during the ongoing starburst as well as low stellar masses, reveals that we start to see SMGs at earlier stages of their evolution.

The improved mapping speed and sensitivity of the new SCUBA2 camera will enable studies of the evolutionary sequence of SMGs using much bigger and more homogeneously selected samples. Moreover, the study of dust production presented here will be pushed forward with a synergy of *Herschel*, SCUBA2, and ultimately ALMA. These facilities will provide a broad wavelength coverage at the IR, which will allow accurate determination of dust temperatures and, in turns, its mass.

We thank Joanna Baradziej, Amy Barger, Rob Ivison and Peter Laursen for discussion and comments; our referee for help with improving this paper; Kirsten Knudsen for kindly providing the data for SMMJ163555.5+661300 prior to publication; and Jorge Iglesias-Páramo for providing his SED templates.

M.J.M. acknowledges support from The Faculty of Science, University of Copenhagen. The Dark Cosmology Centre is funded by the Danish National Research Foundation.

APPENDIX

DUST YIELD CALCULATION

We calculated the dust yield per star required to explain dust mass in a galaxy in the following way. In an IMF with $M_{\min} = 0.15$, $M_{\max} = 120 M_{\odot}$, and a slope $\alpha = 2.35$ (Salpeter 1955, or $\alpha = 1.5$ for top-heavy IMF), the number

of stars with masses between M_0 and M_1 in the stellar population with a total mass of M_* can be expressed as

$$N(M_0 < M < M_1) = M_* \frac{\int_{M_0}^{M_1} M^{-\alpha} dM}{\int_{M_{\min}}^{M_{\max}} M^{-\alpha} M dM}, \quad (\text{A1})$$

where the denominator provides a normalization so that a total mass is equal to M_* . For SNe we assumed $M_0 = 8 M_\odot$ and $M_1 = M_{\max} = 120 M_\odot$, whereas for AGB stars: $M_0 = 2.5 M_\odot$ and $M_1 = 8 M_\odot$.

The average dust yield per star is equal to the dust mass divided by the number of stars, $N(M_0 < M < M_1)$.

REFERENCES

- Alexander D.M., Bauer F.E., Chapman S.C., Smail I., Blain A.W., Brandt W.N., Ivison R.J., 2005, *ApJ*, 632, 736
 Appleton P.N., et al., 2004, *ApJS*, 154, 147
 Archibald E.N., Dunlop J.S., Hughes D.H., Rawlings S., Eales S.A., Ivison R.J., 2001, *MNRAS*, 323, 417
 Barger A.J., Cowie L.L., Richards E.A., 2000, *AJ*, 119, 2092
 Baugh C.M., Lacey C.G., Frenk C.S., Granato G.L., Silva L., Bressan A., Benson A.J., Cole S., 2005, *MNRAS*, 356, 1191
 Beelen A., Cox P., Benford D.J., Dowell C.D., Kovács A., Bertoldi F., Omont A., Carilli C.L., 2006, *ApJ*, 642, 694
 Bell E.F., 2003, *ApJ*, 586, 794
 Benford D.J., Cox P., Omont A., Phillips T.G., McMahon R.G., 1999, *ApJ*, 518, L65
 Bertoldi F., Carilli C.L., Cox P., Fan X., Strauss M.A., Beelen A., Omont A., Zylka R., 2003a, *A&A*, 406, L55
 Bertoldi F., et al., 2003b, *A&A*, 409, L47
 Beswick R.J., Muxlow T.W.B., Thrall H., Richards A.M.S., Garrington S.T., 2008, *MNRAS*, 385, 1143
 Bianchi S., Schneider R., 2007, *MNRAS*, 378, 973
 Blain A.W., Chapman S.C., Smail I., Ivison R., 2004, *ApJ*, 611, 52
 Blair W.P., Ghavamian P., Long K.S., Williams B.J., Borkowski K.J., Reynolds S.P., Sankrit R., 2007, *ApJ*, 662, 998
 Borkowski K.J., et al., 2006, *ApJ*, 642, L141
 Borys C., Chapman S., Halpern M., Scott D., 2003, *MNRAS*, 344, 385
 Boyle B.J., Cornwell T.J., Middelberg E., Norris R.P., Appleton P.N., Smail I., 2007, *MNRAS*, 376, 1182
 Capak P., et al., 2004, *AJ*, 127, 180
 Capak P., et al., 2008, *ApJ*, 681, L53
 Carilli C.L., et al., 2004, *AJ*, 128, 997
 Casey C.M., et al., 2009, *MNRAS*, 400, 670
 Castro Cerón J.M., Michałowski M., Hjorth J., Watson D.J., Fynbo J.P.U., Gorosabel J., 2006, *ApJL*, 653, L85
 Castro Cerón J.M., Michałowski M.J., Hjorth J., Malesani D., Gorosabel J., Watson D., Fynbo J.P.U., 2009, *ApJ*, submitted, [arXiv:0803.2235v1 \[astro-ph\]](#)
 Chabrier G., 2003, *ApJ*, 586, L133
 Chapin E.L., et al., 2009, *MNRAS*, 398, 1793
 Chapman S.C., Richards E.A., Lewis G.F., Wilson G., Barger A.J., 2001, *ApJ*, 548, L147
 Chapman S.C., Blain A.W., Smail I., Ivison R.J., 2005, *ApJ*, 622, 772
 Condon J.J., 1992, *ARA&A*, 30, 575
 Coppin K.E.K., et al., 2009, *MNRAS*, 395, 1905
 Courty S., Björnsson G., Gudmundsson E.H., 2007, *MNRAS*, 376, 1375
 Cowie L.L., Barger A.J., Wang W., Williams J.P., 2009, *ApJ*, 697, L122
 Cox P., et al., 2002, *A&A*, 387, 406
 Daddi E., Dannerbauer H., Krips M., Walter F., Dickinson M., Elbaz D., Morrison G.E., 2009a, *ApJ*, 695, L176
 Daddi E., et al., 2009b, *ApJ*, 694, 1517
 Dannerbauer H., Walter F., Morrison G., 2008, *ApJ*, 673, L127
 Davé R., Finlator K., Oppenheimer B.D., Fardal M., Katz N., Kereš D., Weinberg D.H., 2009, *MNRAS*, submitted, [arXiv:0909.4078 \[astro-ph.CO\]](#)
 Devereux N.A., Young J.S., 1990, *ApJ*, 359, 42
 Devlin M.J., et al., 2009, *Nature*, 458, 737
 Draine B.T., 2003, *ARA&A*, 41, 241
 Draine B.T., 2009, in “Cosmic Dust – Near and Far”, ed. Th. Henning, E. Grün, J. Steinacker (ASP Conf. Series), [arXiv:0903.1658](#)
 Drory N., Bender R., Feulner G., Hopp U., Maraston C., Snigula J., Hill G.J., 2004, *ApJ*, 608, 742
 Dunlop J.S., Hughes D.H., Rawlings S., Eales S.A., Ward M.J., 1994, *Nature*, 370, 347
 Dunne L., Eales S., Ivison R., Morgan H., Edmunds M., 2003, *Nature*, 424, 285
 Dunne L., et al., 2009, *MNRAS*, 394, 1307
 Dwek E., 2004, *ApJ*, 607, 848
 Dwek E., Galliano F., Jones A.P., 2007, *ApJ*, 662, 927
 Elíasdóttir Á., et al., 2007, [arXiv:0710.5636 \[astro-ph\]](#)
 Erb D.K., Steidel C.C., Shapley A.E., Pettini M., Reddy N.A., Adelberger K.L., 2006, *ApJ*, 646, 107
 Ercolano B., Barlow M.J., Sugerman B.E.K., 2007, *MNRAS*, 375, 753
 Ferrarotti A.S., Gail H.P., 2006, *A&A*, 447, 553
 Fontanot F., Monaco P., Silva L., Grazian A., 2007, *MNRAS*, 382, 903
 Frayer D.T., et al., 2006, *ApJ*, 647, L9
 Garn T., Green D.A., Riley J.M., Alexander P., 2009, *MNRAS*, 397, 1101
 Garrett M.A., 2002, *A&A*, 384, L19
 Gehrz R., 1989, In: L. J. Allamandola & A. G. G. M. Tielens (ed.) *Interstellar Dust*, vol. 135 of IAU Symposium, 445
 Gialalisco M., et al., 2004, *ApJ*, 600, L93
 Gomez H.L., Dunne L., Eales S.A., Gomez E.L., Edmunds M.G., 2005, *MNRAS*, 361, 1012
 Gomez H.L., et al., 2009, *MNRAS*, 397, 1621
 Gordon K.D., Clayton G.C., Misselt K.A., Landolt A.U., Wolff M.J., 2003, *ApJ*, 594, 279
 Green D.A., Tuffs R.J., Popescu C.C., 2004, *MNRAS*, 355, 1315
 Greve T.R., et al., 2005, *MNRAS*, 359, 1165
 Greve T.R., Pope A., Scott D., Ivison R.J., Borys C., Conselice C.J., Bertoldi F., 2008, *MNRAS*, 389, 1489
 Greve T.R., et al., 2009, *ApJ*, submitted, [arXiv:0904.0028 \[astro-ph.CO\]](#)
 Gruppioni C., Pozzi F., Zamorani G., Ciliegi P., Lari C., Calabrese E., La Franca F., Matute I., 2003, *MNRAS*, 341, L1
 Hainline L.J., 2008, *Multi-Wavelength Properties of Submillimeter-Selected Galaxies*, Ph.D. thesis, California Institute of Technology
 Helou G., Soifer B.T., Rowan-Robinson M., 1985, *ApJ*, 298, L7
 Hirashita H., 2000, *PASJ*, 52, 585
 Hughes D.H., et al., 1998, *Nature*, 394, 241
 Ibar E., et al., 2008, *MNRAS*, 386, 953
 Iglesias-Páramo J., et al., 2007, *ApJ*, 670, 279
 Iono D., et al., 2006, *ApJ*, 640, L1
 Isaak K.G., Priddey R.S., McMahon R.G., Omont A., Peroux C., Sharp R.G., Withington S., 2002, *MNRAS*, 329, 149
 Ivison R.J., et al., 2010, *MNRAS*, 402, 245
 Kennicutt R.C., 1998, *ARA&A*, 36, 189
 Kippenhahn R., Weigert A., 1990, *Stellar Structure and Evolution*, (Berlin:Springer)
 Knudsen K.K., et al., 2006, *MNRAS*, 368, 487
 Knudsen K.K., Kneib J.P., Egami E., 2008, In: Chary R.R., Teplitz H.I., Sheth K. (eds.) *Infrared Diagnostics of Galaxy Evolution*, vol. 381 of Astronomical Society of the Pacific Conference Series, 372
 Knudsen K.K., Kneib J., Richard J., Petitpas G., Egami E., 2010, *ApJ*, 709, 210
 Kotak R., et al., 2009, *ApJ*, 704, 306
 Kovács A., Chapman S.C., Dowell C.D., Blain A.W., Ivison R.J., Smail I., Phillips T.G., 2006, *ApJ*, 650, 592

- Krause O., Birkmann S.M., Rieke G.H., Lemke D., Klaas U., Hines D.C., Gordon K.D., 2004, *Nature*, 432, 596
- Labbé I., et al., 2005, *ApJ*, 624, L81
- Lacki B.C., Thompson T.A., 2009, *ApJ*, submitted, [arXiv:0910.0478 \[astro-ph.CO\]](#)
- Lacki B.C., Thompson T.A., Quataert E., 2009, *ApJ*, submitted, [arXiv:0907.4161 \[astro-ph.CO\]](#)
- Maiolino R., Schneider R., Oliva E., Bianchi S., Ferrara A., Mannucci F., Pedani M., Roca Sogorb M., 2004, *Nature*, 431, 533
- Marleau F.R., Fadda D., Appleton P.N., Noriega-Crespo A., Im M., Clancy D., 2007, *ApJ*, 663, 218
- Martínez-Sansigre A., et al., 2009, *ApJ*, 706, 184
- Meikle W.P.S., et al., 2007, *ApJ*, 665, 608
- Michałowski M.J., Hjorth J., Castro Cerón J.M., Watson D., 2008, *ApJ*, 672, 817
- Michałowski M.J., et al., 2009, *ApJ*, 693, 347
- Michałowski M.J., Hjorth J., Watson D., 2010, *A&A* accepted, [arXiv:0905.4499](#)
- Morgan H.L., Edmunds M.G., 2003, *MNRAS*, 343, 427
- Morgan H.L., Dunne L., Eales S.A., Ivison R.J., Edmunds M.G., 2003, *ApJ*, 597, L33
- Murphy E.J., 2009, *ApJ*, 706, 482
- Murphy E.J., Chary R.R., Alexander D.M., Dickinson M., Magnelli B., Morrison G., Pope A., Teplitz H.I., 2009, *ApJ*, 698, 1380
- Nozawa T., Kozasa T., Umeda H., Maeda K., Nomoto K., 2003, *ApJ*, 598, 785
- Omont A., Cox P., Bertoldi F., McMahon R.G., Carilli C., Isaak K.G., 2001, *A&A*, 374, 371
- Perera T.A., et al., 2008, *MNRAS*, 391, 1227
- Perley D.A., et al., 2009, *MNRAS*, submitted, [arXiv:0912.2999](#)
- Pope A., et al., 2006, *MNRAS*, 370, 1185
- Portinari L., Sommer-Larsen J., Tantaló R., 2004, *MNRAS*, 347, 691
- Priddey R.S., McMahon R.G., 2001, *MNRAS*, 324, L17
- Priddey R.S., Isaak K.G., McMahon R.G., Robson E.I., Pearson C.P., 2003, *MNRAS*, 344, L74
- Priddey R.S., Ivison R.J., Isaak K.G., 2008, *MNRAS*, 383, 289
- Rho J., et al., 2008, *ApJ*, 673, 271
- Rho J., Reach W.T., Tappe A., Hwang U., Slavin J.D., Kozasa T., Dunne L., 2009, *ApJ*, 700, 579
- Rieke G.H., Alonso-Herrero A., Weiner B.J., Pérez-González P.G., Blaylock M., Donley J.L., Marcellac D., 2009, *ApJ*, 692, 556
- Robson I., Priddey R.S., Isaak K.G., McMahon R.G., 2004, *MNRAS*, 351, L29
- Sajina A., et al., 2008, *ApJ*, 683, 659
- Sakon I., et al., 2009, *ApJ*, 692, 546
- Salpeter E.E., 1955, *ApJ*, 121, 161
- Sandstrom K.M., Bolatto A.D., Stanimirović S., van Loon J.T., Smith J.D.T., 2009, *ApJ*, 696, 2138
- Sargent M.T., et al., 2010, *ApJS*, 186, 341
- Savaglio S., Glazebrook K., LeBorgne D., 2009, *ApJ*, 691, 182
- Scalo J., 1998, In: Gilmore G., Howell D. (eds.) *The Stellar Initial Mass Function* (38th Herstmonceux Conference), vol. 142 of *Astronomical Society of the Pacific Conference Series*, 201
- Schinnerer E., et al., 2008, *ApJ*, 689, L5
- Scott K.S., et al., 2008, *MNRAS*, 385, 2225
- Seaquist E., Yao L., Dunne L., Cameron H., 2004, *MNRAS*, 349, 1428
- Serjeant S., et al., 2003, *MNRAS*, 344, 887
- Seymour N., Huynh M., Dwelly T., Symeonidis M., Hopkins A., McHardy I.M., Page M.J., Rieke G., 2009, *MNRAS*, 398, 1573
- Sibthorpe B., et al., 2009, *ApJ*, submitted [arXiv:0910.1094v1 \[astro-ph.HE\]](#)
- Silva L., Granato G.L., Bressan A., Danese L., 1998, *ApJ*, 509, 103
- Sloan G.C., et al., 2009, *Science*, 323, 353
- Sodroski T.J., Odegard N., Arendt R.G., Dwek E., Weiland J.L., Hauser M.G., Kelsall T., 1997, *ApJ*, 480, 173
- Stark D.P., Bunker A.J., Ellis R.S., Eyles L.P., Lacy M., 2007, *ApJ*, 659, 84
- Sternberg A., 1998, *ApJ*, 506, 721
- Stevens J.A., Amure M., Gear W.K., 2005, *MNRAS*, 357, 361
- Stratta G., Maiolino R., Fiore F., D'Elia V., 2007, *ApJ*, 661, L9
- Sugerman B.E.K., et al., 2006, *Science*, 313, 196
- Tacconi L.J., et al., 2006, *ApJ*, 640, 228
- Tacconi L.J., et al., 2008, *ApJ*, 680, 246
- Todini P., Ferrara A., 2001, *MNRAS*, 325, 726
- Valiante R., Schneider R., Bianchi S., Andersen A.C., 2009, *MNRAS*, 397, 1661
- van der Wel A., Franx M., Wuyts S., van Dokkum P.G., Huang J., Rix H.W., Illingworth G.D., 2006, *ApJ*, 652, 97
- Vlahakis C., Eales S., Dunne L., 2007, *MNRAS*, 379, 1042
- Wang R., et al., 2008, *AJ*, 135, 1201
- Wang W.H., Cowie L.L., Barger A.J., 2004, *ApJ*, 613, 655
- Wang W.H., Cowie L.L., van Saders J., Barger A.J., Williams J.P., 2007, *ApJ*, 670, L89
- Wang W.H., Barger A.J., Cowie L.L., 2009, *ApJ*, 690, 319
- Weiß A., et al., 2009, *ApJ*, 707, 1201
- Wesson R., et al., 2009, *MNRAS*, accepted, [arXiv:0907.0246v1 \[astro-ph.GA\]](#)
- Wilson C.D., et al., 2008, *ApJS*, 178, 189
- Wilson T.L., Batrla W., 2005, *A&A*, 430, 561
- Yang M., Greve T.R., Dowell C.D., Borys C., 2007, *ApJ*, 660, 1198
- Younger J.D., et al., 2007, *ApJ*, 671, 1531
- Younger J.D., et al., 2008a, *MNRAS*, 387, 707
- Younger J.D., et al., 2008b, *ApJ*, 688, 59
- Younger J.D., et al., 2009a, *ApJ*, 704, 803
- Younger J.D., et al., 2009b, *MNRAS*, 394, 1685
- Zafar T., et al., 2010, *A&A*, accepted, [arXiv:1002.4101](#)
- Zhukovska S., Gail H.P., Tieloff M., 2008, *A&A*, 479, 453